



**INFLUENZA UPDATE  
2011-2012  
DEBBYE ROSEN  
ADULT IMMUNIZATION COORDINATOR  
CONNECTICUT DEPARTMENT OF PUBLIC HEALTH  
860-509-7729 [debbye.rosen@ct.gov](mailto:debbye.rosen@ct.gov)**

# 2011-2012 RECOMMENDATIONS

*Centers for Disease Control and Prevention*

# MMWR

Morbidity and Mortality Weekly Report

Early Release / Vol. 60

August 18, 2011

**Prevention and Control of Influenza with Vaccines: Recommendations of the  
Advisory Committee on Immunization Practices (ACIP), 2011**

**EARLY RELEASE AUGUST 18, 2011**

**<http://www.cdc.gov/mmwr/>**



# 2011-2012 RECOMMENDATIONS

Centers for Disease Control and Prevention

# MMWR

Morbidity and Mortality Weekly Report

Weekly / Vol. 60 / No. 33

August 26, 2011

1128 Prevention and Control of Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices (ACIP), 2011

1127 Notes from the Field: Mortality Among Refugees



<http://www.cdc.gov/mmwr/PDF/wk/mm6033.pdf>


# 2011-2012 ACIP

- **Vaccination of all persons aged  $\geq 6$  months (adopted in 2010) continues to be recommended.**
- **Vaccine virus strains for 2011–2012**
- **The vaccination schedule for children aged 6 months through 8 years**
- **Considerations regarding vaccination of persons with egg allergy**
- **A new formulation of trivalent inactivated vaccine that is given intradermally (into the skin)**

2011-2012

## WHO YOU SHOULD VACCINATE

- **160 MILLION DOSES WILL BE AVAILABLE**
- **≥ 6 MONTHS OF AGE**
- **AT RISK POPULATIONS**
  - HEART DISEASE, COPD AND OTHER CHRONIC RESPIRATORY ILLNESSES, DIABETES**
- **PREGNANT**
- **VACCINATE AS SOON AS VACCINE IS AVAILABLE**



# 2011-2012



## Vaccine strains for 2011-2012


- A/California/7/2009-like H1N1
- A/Perth/16/2009-like H3N2
- B/Brisbane/60/2008

# CHILDREN 6 MONTHS - 8 YEARS




- **CHILDREN IN THIS AGE GROUP RECEIVING VACCINE FOR THE FIRST TIME OR WITH UNKNOWN VACCINE HISTORY SHOULD RECEIVE 2 DOSES**
- **CHILDREN IN THIS AGE GROUP WHO RECEIVED AT LEAST 1 DOSE OF 2010-2011 SEASONAL VACCINE WILL REQUIRE ONLY ONE DOSE OF 2011-2012 SEASONAL VACCINE**
- **CHILDREN IN THIS AGE GROUP WHO DID NOT RECEIVE AT LEAST ONE DOSE OF 2010-2011 VACCINE OR FOR WHOM IT IS NOT CERTAIN WHETHER THE 2010-2011 DOSE WAS RECEIVED, SHOULD RECEIVE 2 DOSES**

# CHILDREN 6 MONTHS- 8 YEARS



# EGG ALLERGY



\* Persons with egg allergy might tolerate egg in baked products (e.g., bread or cake). Tolerance to egg-containing foods does not exclude the possibility of egg allergy.

## 2011-2012 INFLUENZA VACCINE PRODUCTS



Product Name	Age Indication	Manufactured by
Fluarix	≥ 3 years of age	GlaxoSmithKline
FluLaval	≥ 18 years of age	ID Biomedical Corporation
Fluvirin	≥ 4 years of age	Novartis
Agriflu	≥ 18 years of age	Novartis
Afluria	> 5 years of age*	CSL Limited (Merck)
Fluzone	> 6 months of age	Sanofi Pasteur
High dose Fluzone	≥ 65 years of age	Sanofi Pasteur
Intradermal Fluzone	18 years-64 years of age	Sanofi Pasteur
FluMist-nasal	2 years-49 years**	MedImmune
*In the US not recommended for children 5-8yrs **healthy		

Package Inserts: Influenza - Windows Internet Explorer

http://www.immunize.org/packageinserts/pi\_influenza.asp

Convert Select

Favorites Department of Public Health Suggested Sites Upgrade Your Browser

Package Inserts: Influenza

Handouts for Patients & Staff Clinic Resources Vaccine Information Statements Diseases & Vaccines Talking about Vaccines Topics

IAC Resources

- IAC Home
- Resources Home
- Handouts
- Clinic Resources
- Shop IAC
- VISs
- IAC Image Library
- IAC Publications
- Adult Vax Guide

Other Resources

- Books and Periodicals
- Coalitions Websites
- Continuing Education
- Email News Services
- Government Agencies
- Hotlines
- International Organizations
- Manufacturers
- Package Inserts
- Partners
- PowerPoint

## Package Inserts

### Vaccine index

<a href="#">View all</a>	<a href="#">HPV</a>	<a href="#">Pneumococcal - PPSV</a>	<a href="#">Td</a>
<a href="#">Anthrax</a>	<a href="#">Influenza</a>	<a href="#">Polio - IPV</a>	<a href="#">Tdap</a>
<a href="#">Chickenpox (varicella)</a>	<a href="#">Japanese encephalitis</a>	<a href="#">Rabies</a>	<a href="#">Typhoid</a>
<a href="#">DTaP</a>	<a href="#">Measles</a>	<a href="#">Rotavirus</a>	<a href="#">Yellow fever</a>
<a href="#">Hib</a>	<a href="#">Meningococcal</a>	<a href="#">Rubella</a>	
<a href="#">Hepatitis A</a>	<a href="#">Mumps</a>	<a href="#">Shingles (zoster)</a>	
<a href="#">Hepatitis B</a>	<a href="#">Pneumococcal - PCV</a>	<a href="#">Smallpox</a>	

### Package Inserts - Influenza

<a href="#">Afluria® Package Insert (injectable)</a> CSL Biotherapies Inc.	<a href="#">Fluvirin® Package Insert (injectable)</a> Novartis Vaccines
<a href="#">Aqriflu Package Insert (injectable)</a> Novartis	<a href="#">Fluzone High-Dose Package Insert (injectable)</a> sanofi pasteur
<a href="#">Fluarix® Package Insert (injectable)</a> GlaxoSmithKline	<a href="#">Fluzone® Package Insert (injectable)</a> sanofi pasteur
<a href="#">FluLaval® Package Insert (injectable)</a> GlaxoSmithKline	<a href="#">FluMist® Package Insert (intranasal)</a> MedImmune, Inc

Internet | Protected Mode: Off

125%

11:11 AM  
8/9/2011



[http://www.immunize.org/packageinserts/pi\\_influenza.asp](http://www.immunize.org/packageinserts/pi_influenza.asp)

## INTRADERMAL INFLUENZA VACCINE

Fluzone Intradermal vaccine is an inactivated influenza virus vaccine indicated for active immunization of persons 18 through 64 years of age against influenza disease caused by influenza virus subtypes A and type B contained in the vaccine.

Please see Important Safety Information on reverse side.



**With a microneedle delivery system that's a mere 1.5mm long, Fluzone Intradermal vaccine is a new choice in influenza immunization.**

**Available for the 2011-2012 influenza season.**

— [actual size]

# INTRADERMAL VACCINE ADMINISTRATION






*School Immunization Requirement Update March 9, 2011*



VACCINE	GRADE	# of doses	Start Date
Pneumococcal	Pre-K and K (born 1/1/2007 or later and less than 5 years old)	1 dose on or after 1 <sup>st</sup> birthday	August 1, 2011
Hepatitis A	Pre-K and K (born 1/1/2007 or later)	2 doses given six months apart - 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday	August 1, 2011
<b>Influenza</b>	<b>Pre-K (children age 24-59 months) given annually between August 1<sup>st</sup> and December 31<sup>st</sup> each year</b>	<b>1 dose - (2 doses for those receiving flu vaccine for 1<sup>st</sup> time)</b>	<b>August 1, 2011</b>
MMR	K-12	2 doses given at least 28 days apart - 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday	August 1, 2011
Varicella	Pre-K	1 dose on or after 1 <sup>st</sup> birthday	August 1, 2011
Varicella	K and 7 <sup>th</sup> grade entry	2 doses given 3 months apart - 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday	August 1, 2011
Tdap	7 <sup>th</sup> grade entry	1 dose	August 1, 2011
Meningococcal	7 <sup>th</sup> grade entry	1 dose	August 1, 2011

*new/ sch/ regs/ provider ltr/ 3/09/2011*



**By January 1, 2011 and each January 1 thereafter, children aged 6–59 months attending a child day care center, group day care home, or family day care home are required to receive at least one dose of influenza vaccine between September 1 and December 31 of the preceding year. If**

## LEARN MORE ABOUT IT



[www.cdc.gov/flu](http://www.cdc.gov/flu)

[www.ct.gov/dph/immunizations](http://www.ct.gov/dph/immunizations)

[http://www.immunize.org/packageinserts/pi\\_influenza.asp](http://www.immunize.org/packageinserts/pi_influenza.asp)

[www.cdc.gov/hepatitis/Partners/PeriHepBCoord.htm](http://www.cdc.gov/hepatitis/Partners/PeriHepBCoord.htm)



**INFLUENZA UPDATE  
2011-2012  
DEBBYE ROSEN  
ADULT IMMUNIZATION COORDINATOR  
CONNECTICUT DEPARTMENT OF PUBLIC HEALTH  
860-509-7729 [debbye.rosen@ct.gov](mailto:debbye.rosen@ct.gov)**

??

